

PACE-IIT & MEDICAL

MUMBAI / AKOLA / DELHI / KOLKATA / GHAZIABAD / NASHIK / GOA / BOKARO / PUNE

IIT – JEE: 2024

TW TEST (MAIN)

DATE: 28/08/22

TOPIC: VECTORS

Answer Key

- | | | | | |
|---------|---------|---------|---------|---------|
| 1. (A) | 2. (C) | 3. (C) | 4. (D) | 5. (B) |
| 6. (A) | 7. (B) | 8. (B) | 9. (D) | 10. (C) |
| 11. (B) | 12. (B) | 13. (C) | 14. (B) | 15. (B) |
| 16. (D) | 17. (D) | 18. (A) | 19. (C) | 20. (C) |
| 21. (D) | 22. (A) | 23. (A) | 24. (D) | 25. (A) |

SOLUTIONS

1. (A)

$$\frac{\vec{b} \cdot \vec{a}}{|\vec{a}|} \hat{a}$$
$$= \frac{8(\hat{i} + 2\hat{j} + 2\hat{k})}{3}$$
$$= \frac{8}{9}(\hat{i} + 2\hat{j} + 2\hat{k})$$

2. (C)

$$R^2 = (3P)^2 + (2P)^2 + 2(3P)(2P)\cos\theta$$
$$\Rightarrow R^2 = 13P^2 + 12P^2 \cos\theta \quad \text{.....(1)}$$
$$(2R)^2 = (6P)^2 + (2P)^2 + 2(6P)(2P)\cos\theta$$
$$4R^2 = 40P^2 + 24P^2 \cos\theta \quad \text{.....(2)}$$

From (1) & (2)

$$\cos\theta = -\frac{1}{2} \Rightarrow \theta = 120^\circ$$

3. (C)

From triangle law

$$\vec{C} + \vec{A} = \vec{B} \quad \text{.....(C)}$$

4. (D)

$$\vec{F}_{\text{net}} = 0$$

$$(-5\hat{i} + 5\hat{j}) + (3\hat{i} + 3\hat{j}) + \vec{F}_3 = 0$$

$$-2\hat{i} + 8\hat{j} + \vec{F}_3 = 0$$

$$\vec{F}_3 = 2\hat{i} - 8\hat{j}$$

5. (B)

$$\vec{A} + \vec{B} + \vec{C} + \vec{D} + \vec{E} + \vec{F} = \vec{0}$$

$$\vec{A} = -\vec{D}$$

6. (A)

$$\vec{A} - \vec{B} = 3\hat{i} + \hat{k}$$

$$\text{Unit vector of } \vec{A} - \vec{B} = \frac{3\hat{i} + \hat{k}}{\sqrt{10}}$$

7. (B)

Sum of any three should be greater than fourth

8. (B)

$$\vec{V}_1 = 50\hat{j} \quad \vec{V}_2 = -50\hat{j}$$

$$\Delta\vec{V} = \vec{V}_2 - \vec{V}_1 = -50\hat{i} - 50\hat{j}$$

$$= 50\sqrt{2}\text{km/h (S - W)}$$

9. (D)

$$\vec{V} = 10 \frac{(3\hat{i} + 4\hat{j})}{5} = 6\hat{i} + 8\hat{j}$$

10. (C)

$$R^2 = P^2 + Q^2 + 2PQ \cos 60^\circ$$

$$7Q^2 = P^2 + Q^2 + 2PQ \times \frac{1}{2}$$

$$\left(\frac{P}{Q}\right)^2 - 6 + \frac{P}{Q} = 0$$

$$\frac{P}{Q} = 2$$

11. (B)

$$\vec{S} = \hat{i} + (-5\hat{j}) + 2\hat{i} + 9\hat{j} = 3\hat{i} + 4\hat{j}$$

$$|\vec{S}| = 5 \text{ miles}$$

12. (B)

$$W = \vec{F} \cdot \vec{S} = -12 + 2c - 6 = 6$$

$$c = 12$$

13. (C)

$$\text{Let } \vec{b} = x\hat{i} + y\hat{j} + z\hat{k}$$

$$\vec{a} \cdot \vec{b} = x + y + z = 1 \quad \text{_____ (1)}$$

$$\vec{a} \times \vec{b} = (z - y)\hat{i} + (x - z)\hat{j} + (y - x)\hat{k} = \hat{j} - \hat{k} \quad \text{_____ (2)}$$

$$y = z = 0 \quad x - z = 1$$

$$y - x = -1$$

$$x = 1$$

$$\therefore \vec{b} = \hat{i}$$

14. (B)

$$\vec{A} + \vec{B} = 3\hat{i} + 7\hat{j} + \hat{k}$$

$$\text{Unit vector } \frac{3\hat{i} + 7\hat{j} + \hat{k}}{\sqrt{59}}$$

15. (B)

$$\vec{A} \times \vec{B} = 8\hat{i} - 8\hat{j} - 8\hat{k}$$

$$|\vec{A} \times \vec{B}| = 8\sqrt{3}$$

16. (D)

$$(\hat{i} + \hat{j}) \cdot (\hat{i} - \hat{j}) = 1 - 1 = 0$$

17. (D)

$$\vec{A} \perp \vec{B} \& \vec{A} \perp \vec{C} \text{ then } \vec{A} \parallel (\vec{B} \times \vec{C})$$

18. (A)

$$\vec{V} = \vec{\omega} \times \vec{r} = -2\hat{i} + 3\hat{j} + 4\hat{k}$$

$$|\vec{V}| = \sqrt{29}$$

19. (C)

$$|\vec{V}_1 + \vec{V}_2|^2 = |\vec{V}_1 - \vec{V}_2|^2$$

$$V_1^2 + V_2^2 + 2V_1V_2 \cos \theta = V_1^2 + V_2^2 - 2V_1V_2 \cos \theta$$

$$4V_1V_2 \cos \theta = 0$$

$$\theta = 90^\circ$$

$$\vec{V}_1 \perp \vec{V}_2$$

20. (C)

Direction cosines

$$\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$$

$$\therefore \sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma = 2$$

21. (D)

$$\overrightarrow{AB} + \overrightarrow{AC} + \overrightarrow{AD} + \overrightarrow{AE} + \overrightarrow{AF}$$

$$\overrightarrow{AB} + (\overrightarrow{AC} + \overrightarrow{AF}) + \overrightarrow{AD} + \overrightarrow{AD} + \overrightarrow{DE}$$

$$\overrightarrow{AF} = \overrightarrow{CD} \text{ \& } \overrightarrow{AB} = -\overrightarrow{DE}$$

$$3\overrightarrow{AD} = 3(2\overrightarrow{AO}) = 6\overrightarrow{AO}$$

22. (A)

$$\frac{\overrightarrow{A} \cdot \overrightarrow{B}}{|\overrightarrow{B}|} \hat{B} = \frac{5}{\sqrt{2}} \left(\frac{\hat{i} + \hat{j}}{\sqrt{2}} \right)$$

$$= \frac{5}{2} (\hat{i} + \hat{j})$$

23. (A)

$$\overrightarrow{A} + \overrightarrow{B} + \overrightarrow{C} = 0$$

$$\overrightarrow{C} = -(\overrightarrow{A} + \overrightarrow{B}) = -(3\hat{i} + 4\hat{k})$$

24. (D)

If $\overrightarrow{A} \cdot \overrightarrow{B} = 0$ then $\overrightarrow{A} \perp \overrightarrow{B}$

25. (A)

$$\overrightarrow{P} \cdot \overrightarrow{A} = 0 \Rightarrow a^2 - 2a - 3 = 0$$

$$(a+1)(a-3) = 0$$

$$a = -1, 3$$

PACE-IIT & MEDICAL

MUMBAI / AKOLA / DELHI / KOLKATA / GHAZIABAD / NASHIK / GOA / BOKARO / PUNE

IIT – JEE: 2024

TW TEST (MAIN)

DATE: 28/08/22

TOPIC: MOLE CONCEPT

Answer Key

26. (C)	27. (A)	28. (C)	29. (C)	30. (A)
31. (C)	32. (C)	33. (C)	34. (D)	35. (C)
36. (C)	37. (D)	38. (D)	39. (B)	40. (B)
41. (D)	42. (A)	43. (D)	44. (D)	45. (B)
46. (D)	47. (A)	48. (D)	49. (C)	50. (B)

SOLUTIONS

26. (C)

27. (A)

28. (C)

29. (C)

30. (A)


31. (C)

32. (C)




$$= 7 + 24 + 1$$

$$= 32$$



$$\text{Mole of NO}_3^- = \frac{3.1 \times 10^{-3}}{62} = \frac{10^{-3}}{20} = \frac{10^{-4}}{2} = 5 \times 10^{-5}$$



1

→

32

$$5 \times 10^{-5} \times 6 \times 10^{23}$$



$$32 \times 30 \times 10^{18}$$

$$= 30 \times 10^{18}$$

$$= 96 \times 10^{19}$$

33. (C)

$$\text{Mole of H}_2\text{O} = \frac{0.45}{18} = \frac{5}{200}$$



$$\frac{5}{200} \quad \frac{5}{200} \times 8$$

$$= \frac{40}{200} = 0.2$$

$$\begin{aligned} \text{No. of neutrons} &= 0.2 \times 6 \times 10^{23} \\ &= 1.2 \times 10^{23} \end{aligned}$$

34. (D)



35. (C)

36. (C)

37. (D)

38. (D)

39. (B)



$$\frac{1.5 \times 4}{3} \quad 1.5$$

= 2 mole

$$\text{Mass of Al} = 2 \times 27 = 54 \text{ gm}$$

40. (B)

41. (D)

42. (A)

43. (D)

44. (D)

- 45. (B)
- 46. (D)
- 47. (A)
- 48. (D)
- 49. (C)
- 50. (B)

PACE-IIT & MEDICAL

MUMBAI / AKOLA / DELHI / KOLKATA / GHAZIABAD / NASHIK / GOA / BOKARO / PUNE

IIT – JEE: 2024

TW TEST (MAIN)

DATE: 28/08/22

TOPIC: QUADRATIC EQUATIONS

Answer Key

51. (D)	52. (A)	53. (C)	54. (D)	55. (A)
56. (A)	57. (C)	58. (D)	59. (D)	60. (B)
61. (B)	62. (D)	63. (D)	64. (B)	65. (B)
66. (A)	67. (C)	68. (C)	69. (D)	70. (B)
71. (B)	72. (D)	73. (D)	74. (C)	75. (D)

SOLUTION

51. (D)

$$A = 1, B = -2(a + b), C = 2(a^2 + b^2)$$

$$B^2 - 4AC = 1[2(a + b)]^2 - 4(1)(2a^2 + 2b^2)$$

$$= 4a^2 + 4b^2 + 8ab - 8a^2 - 8b^2$$

$$= -4a^2 - 4b^2 + 8ab$$

$$= -4(a - b)^2 < 0$$

So roots are imaginary and different.

52. (A)

The required equation is

$$x^2 - \left\{ (2 + \sqrt{3}) + (2 - \sqrt{3}) \right\} x + (2 + \sqrt{3})(2 - \sqrt{3}) = 0$$

$$\text{Or } x^2 - 4x + 1 = 0$$

53. (C)

Roots are of opposite sign so


$$f(0) < 0$$

$$\lambda^2 - 5\lambda + 6 < 0$$

$$(\lambda - 2)(\lambda - 3) < 0$$

$$\Rightarrow \lambda \in (2, 3)$$

54. (D)



$$f(1) > 0$$

$$f(2) < 0$$

$$f(3) > 0$$

$$\text{So } 1 + \lambda + 1 + \lambda^2 - 3\lambda - 6 > 0 \Rightarrow \lambda^2 - 2\lambda - 4 > 0$$

$$\lambda \in (-\infty, 1 - \sqrt{5}) \cup (1 + \sqrt{5}, \infty)$$

$$f(2) = 4 + 2\lambda + 2 + \lambda^2 - 3\lambda - 6 < 0 \Rightarrow \lambda^2 - \lambda < 0$$


$$\lambda(\lambda - 1) < 0$$

$$\lambda \in (0, 1)$$

$$f(3) = 9 + 3\lambda + 3 + \lambda^2 - 3\lambda - 6 > 0$$

$$\lambda^2 + 6 > 0$$

$$\lambda \in \mathbb{R}$$



55. (A)

α, β are roots of equation $x^2 + px + q = 0$ so

$$\alpha + \beta = -p$$

$$\alpha\beta = q$$

And $\alpha^2 + p\alpha + q = 0$

$$\alpha + p = \frac{-q}{\alpha}$$

$$\text{So } (\alpha + p)^{-2} = \frac{\alpha^2}{q^2} \quad \text{similarly } (\beta + p)^{-2} = \frac{\beta^2}{q^2}$$

So required equation

$$x^2 - \left(\frac{\alpha^2 + \beta^2}{q^2} \right) x + \frac{\alpha^2 \beta^2}{q^4} = 0 \Rightarrow x^2 - \left(\frac{p^2 - 2q}{q^2} \right) x + \frac{1}{q^2} = 0 \Rightarrow q^2 x^2 - (p^2 - 2q)x + 1 = 0$$

56. (A)

One root of equation $x^2 + Ax + 12 = 0$ is 4.

$$\text{Sum of roots} = -A$$

$$\text{Product of roots} = 12$$

So other root is 3.

$$A = -7$$

Now roots of equation $x^2 + 2Ax + B = 0$ are equal

$$\text{So } 4A^2 - 4B = 0$$

$$B = A^2 \Rightarrow B = 49$$

57. (C)


Roots are real so $4q^2 - 4pr \geq 0$

And $4pr - 4q^2 \geq 0$

Both inequations are true if

$$q^2 - pr = 0 \Rightarrow \frac{p}{q} = \frac{q}{r}$$

58. (D)




$$f(x) = 2x^2 + \lambda x - (\lambda + 1) = 0$$

$$D = \lambda^2 + 8(\lambda + 1) \geq 0$$

$$\lambda \in [-\infty, -4 - 2\sqrt{2}] \cup [-4 + 2\sqrt{2}, \infty] \quad \dots\dots\dots (1)$$

$$\frac{-b}{2a} = \frac{-\lambda}{4} > 1 \Rightarrow \lambda < -4 \quad \dots\dots\dots (2)$$

$$f(1) = 2 + \lambda - (\lambda + 1) > 0, \forall \lambda \in \mathbb{R}$$



$$\Rightarrow \lambda \in (-\infty, -4 - 2\sqrt{2})$$

59. (D)

It is identity in x so

$$r^2 - 2r + 1 = 0 \Rightarrow r = 1$$

$$r^2 - 3r + 2 = 0 \Rightarrow r = 1, 2$$

$$r^2 + 4r + 3 = 0 \Rightarrow r = -1, -3$$

No common value of r is possible so it is not possible.

60. (B)

$2x^2 - 5x - 7 = 0$ roots are α, β

$$\text{So } \alpha + \beta = \frac{5}{2}$$

$$\alpha\beta = -\frac{7}{2}$$

Now roots are $\frac{\alpha}{\beta}, \frac{\beta}{\alpha}$

$$\text{Sum} = \frac{\alpha^2 + \beta^2}{\alpha\beta} = \frac{(\alpha + \beta)^2 - 2\alpha\beta}{\alpha\beta} = \frac{\frac{25}{4} + 7}{\frac{-7}{2}} = -\frac{53}{14}$$

Product = 1

So equation is $x^2 + \frac{53}{14}x + 1 = 0$

$$14x^2 + 53x + 14 = 0$$

61. (B)

$ax^2 + bx + c = 0$ roots are imaginary so $b^2 - 4ac < 0$

$$(4c + 2b + a)x^2 - 2(a + b)x + a = 0$$

$$D = 4(a + b)^2 - 4a(4c + 2b + a)$$

$$= 4[a^2 + b^2 + 2ab - 4ac - 2ab - a^2]$$

$$= 4(b^2 - 4ac) < 0$$

So roots are imaginary.

62. (D)

α, β, γ are root of $x^3 - 5x^2 + x - 2 = 0$

$$\text{Now } y = \frac{\alpha + 2}{\alpha - 2} \Rightarrow \alpha = \frac{2(y + 1)}{y - 1}$$

It is root of given equation so

$$\frac{8(y + 1)^3}{(y - 1)^3} - 5.4 \frac{(y + 1)^2}{(y - 1)^2} + \frac{2(y + 1)}{(y - 1)} - 2 = 0$$

$$8(y + 1)^3 - 20(y + 1)^2(y - 1) + 2(y + 1)(y - 1)^2 - 2(y - 1)^3 = 0 \Rightarrow 3y^3 - 2y^2 - 9y - 8 = 0$$

Roots of this equation are $\frac{\alpha + 2}{\alpha - 2}, \frac{\beta + 2}{\beta - 2}, \frac{\gamma + 2}{\gamma - 2}$

So product of roots

$$\left(\frac{\alpha + 2}{\alpha - 2}\right)\left(\frac{\beta + 2}{\beta - 2}\right)\left(\frac{\gamma + 2}{\gamma - 2}\right) = \frac{8}{3}$$

63. (D)

Equation $8x^3 + 1001x + 2008 = 0$ has roots r, s and t .

$$r + s + t = 0, rst = -\frac{2008}{8} = -251$$

Now, let $r + s = A, s + t = B, t + r = C$.

$$\therefore A + B + C = 2(r + s + t) = 0$$

Hence,

$$A^3 + B^3 + C^3 = 3ABC$$

$$\therefore (r + s)^3 + (s + t)^3 + (t + r)^3$$

$$\begin{aligned}
&= 3(r+s)(s+t)(t+r) \\
&= 3(r+s+t-t)(s+t+r-r)(t+r+s-s) \\
&= 3(251) = 753
\end{aligned}$$

64. (B)

Let,

$$\frac{x}{x^2 - 5x + 9} = y$$

$$\Rightarrow yx^2 - 5yx + 9y = x$$

$$\Rightarrow yx^2 - (5y+1)x + 9y = 0$$

Now, x is real, so

$$D \geq 0$$

$$\Rightarrow (-(5y+1))^2 - 4y(9y) \geq 0$$

$$\Rightarrow -11y^2 + 10y + 1 \geq 0$$

$$\Rightarrow 11y^2 - 10y - 1 \leq 0$$

$$\Rightarrow (11y+1)(y-1) \leq 0$$

$$\Rightarrow -\frac{1}{11} \leq y \leq 1$$

65. (B)

Let,

$$y = \frac{3x^2 + 9x + 17}{3x^2 + 9x + 7}$$

$$\Rightarrow 3x^2(y-1) + 9x(y-1) + 7y - 17 = 0$$

Since x is real, so,

$$D \geq 0$$

$$\Rightarrow 81(y-1)^2 - 4 \times 3(y-1)(7y-17) \geq 0$$

$$\Rightarrow (y-1)(y-41) \leq 0 \Rightarrow 1 \leq y \leq 41$$

Therefore, the maximum value of y is 41.

66. (A)

$$D = b^2 - 4a < 0 \Rightarrow a > 0$$

Therefore the graph is concave upwards.

$$f(x) > 0, \forall x \in \mathbb{R}$$

$$\Rightarrow f(-1) > 0$$

$$\Rightarrow a + b + 1 > 0$$

67. (C)

$$(\alpha - \gamma)(\alpha - \delta) = \alpha^2 - (\gamma + \delta)\alpha + \gamma\delta$$

$$= \alpha^2 + p\alpha - 4 \quad (\text{because } \gamma + \delta = -p, \gamma\delta = -4)$$

α is root of $x^2 + px + 7 = 0$

$\Rightarrow \alpha^2 + p\alpha + 7 = 0 \Rightarrow \alpha^2 + p\alpha = -7$

Now $(\alpha - \gamma)(\alpha - \delta) = \alpha^2 + p\alpha - 4 = -7 - 4 = -7 - 4 = -11$

68. (C)

$2x^2 + 9x + 4a = 0$ one root is α then 2α is root at $2x^2 + 3x + a = 0$

So $2\alpha^2 + 9\alpha + 4a = 0$ (1)

$8\alpha^2 + 6\alpha + a = 0$ (2)

By solving (1) and (2) $\alpha = -\frac{a}{2}$

From (1)


$2 \cdot \frac{a^2}{4} - \frac{9a}{2} + 4a = 0$ So $a = 1$

69. (D)

$y(-4) = 16 - 24 - 4 = -12$ min. value = $\frac{-D}{4a} = \frac{-(36+16)}{4} = -13$

$y(3) = 9 + 18 - 4 = 23$

So range is $[-13, 23)$



70. (B)

Correct equation

$x^2 + 13x + q = 0$ (1)

Incorrect equation is

$x^2 + 17x + q = 0$ (2)

Given that roots of Eq. (1) are -2 and -15 .

Therefore, product of the roots of incorrect equation is $q = (-2)(-15) = 30$. From (1), the correct equation is

$x^2 + 13x + 30 = 0$

$\therefore x = -3, -10$

71. (B)

Trick: By inspection, we see that all the values of x lying in $(-\infty, 1) \cup (2, 3)$ satisfy the equations and no other value outside the interval satisfy it.

72. (D)

8, 2 are the roots of $x^2 + ax + \beta = 0$,

$\therefore 8+2=10=-a$, $8 \cdot 2=16=\beta$, i.e. $a=-10$, $\beta=16$ 3, 3 are the roots of

$$x^2 + \alpha x + b = 0 \quad \therefore 3+3=6=-\alpha$$

i.e. $\alpha=-6$, $b=9$

Now, $x^2 + ax + b = 0$ becomes $x^2 - 10x + 9 = 0$ or $(x-1)(x-9) = 0 \Rightarrow x=1, 9$

73. (D)

Here $ax^2 + bx + c = a(x-\alpha)(x-\beta)$

Since, α, β be the roots of $ax^2 + bx + c = 0$.

Also, $\alpha < k < \beta$, so $a(k-\alpha)(k-\beta) < 0$

Also, $a^2k^2 + abk + ac = a(ak^2 + bk + c) = a^2(k-\alpha)(k-\beta) < 0 \Rightarrow a^2k^2 + abk + ac < 0$

74. (C)

$$k-2 > 0 \quad k > 2 \text{ and}$$

$$64 - 4(k+4)(k-2) < 0$$

$$16 - (k^2 + 2k - 8) < 0$$

$$16 - k^2 - 2k + 8 < 0$$

$$-k^2 - 2k + 24 < 0$$

$$k^2 + 2k - 24 > 0$$

$$(k-4)(k+6) > 0 \Rightarrow k \in (-\infty, -6) \cup (4, \infty)$$

Hence, $k \in (4, \infty)$

75. (D)

$$\alpha + \beta + \gamma = 0,$$

$$\frac{\alpha^3 - 3\alpha + \beta^3 - 3\beta + \gamma^3 - 3\gamma}{\alpha\beta\gamma} = \frac{(\alpha^3 + \beta^3 + \gamma^3) - 3(\alpha + \beta + \gamma)}{\alpha\beta\gamma}$$

$$= \frac{3\alpha\beta\gamma - 0}{\alpha\beta\gamma} = 3$$